Species Tag:	19001	Name:	18OH
Version:	3		Hydroxyl radical
Date:	June 2013		X ${}^{2}\Pi$ states
Contributor:	B. J. Drouin		v = 0,1,2
Lines Listed: Freq. (GHz) < Max. J: LOGSTR0= LOGSTR1= Isotope Corr.: Egy. (cm <sup>-1</sup> ) > $\mu_a = \mu_b = \mu_c = \mu_c = \mu_c$	3159 70000 50 -30.0 -35.0 0.0 0.0 1.65520	Q(300.0) = Q(225.0) = Q(150.0) = Q(75.00) = Q(37.50) = Q(18.75) = Q(9.375) = A = B = C =	40.32992 22.82372 17.05378

The positions and energies are determined from a multi-isotopic analysis of ground and v < 2 data as decribed in B.J. Drouin.JCPA 2012, DOI: 10.1021/jp400923z and references therein.

The dipole moment is from K. I. Peterson, G. T. Fraser, and W. A. Klemperer, 1984, Can. J. Phys. **62**, 1502.

state label	vibrational quantum number	isotopologue
0	0	$^{16}{\rm O}^{1}{\rm H}$
1	1	$^{16}\mathrm{O}^{1}\mathrm{H}$
2	2	$^{16}{\rm O}^{1}{\rm H}$
10	0	$^{17}\mathrm{O}^{1}\mathrm{H}$
11	1	$^{17}\mathrm{O}^{1}\mathrm{H}$
12	2	$^{17}\mathrm{O}^{1}\mathrm{H}$
20	0	$^{18}\mathrm{O}^{1}\mathrm{H}$
21	1	$^{18}\mathrm{O}^{1}\mathrm{H}$
22	2	$^{18}\mathrm{O}^{1}\mathrm{H}$
30	0	$^{16}\mathrm{O}^2\mathrm{H}$
31	1	$^{16}\mathrm{O}^2\mathrm{H}$
32	2	$^{16}\mathrm{O}^2\mathrm{H}$
40	0	$^{17}\mathrm{O}^2\mathrm{H}$
41	1	$^{17}\mathrm{O}^{2}\mathrm{H}$
42	2	$^{17}\mathrm{O}^2\mathrm{H}$
50	0	$^{18}{ m O}^{2}{ m H}$
51	1	$^{18}\mathrm{O}^{2}\mathrm{H}$
52	2	$^{18}\mathrm{O}^{2}\mathrm{H}$