

Species Tag:	36005	Name:	HCl ⁺
Version:	2		Chloroniumyl cation
Date:	Dec. 2016		v = 1 ← 0
Contributors:	B. J. Drouin		² Π _{1/2} ← ² Π _{3/2}
			² Σ ← ² Π
Lines Listed:	2583	Q(300.0)=	196.9653
Freq. (GHz) <	86000	Q(225.0)=	147.6317
Max. J:	40	Q(150.0)=	102.4779
LOGSTR0=	-8.0	Q(75.00)=	59.9414
LOGSTR1=	-10.0	Q(37.50)=	40.3231
Isotope Corr.:	-0.122	Q(18.75)=	33.1500
Egy. (cm ⁻¹) >	0.0	Q(9.375)=	31.9340
μ ₀ =	1.0	A=	
μ' =	0.1959	B=	293443.75
μ _{el} =	0.13	C=	

The work of H. Gupta, B. J. Drouin, & J. C. Pearson, 2012, ApJ, **751**, L38 and the optical spectra in W. D. Sheasley, 1972, Ph.D. Dissertation, The Ohio State University; Ann Arbor, MI. is expanded to include vibrational data from Doménech, Drouin, Cernicharo *et al.* ApJL 833 L32 (2016). The transition dipole moments for infrared (μ' and optical (μ_{el}) are calculated by A. Pradhan *et al.* 1991, J. Chem. Phys, **95**, 9010. The magnetic dipole, which predicts fine structure transitions in the mid and far-ir was taken to be 1 Bohr magneton. The state identifiers v = 80 and v = 81 refer to the ground and first excited vibrational levels, respectively. The ²Σ ground vibrational state has a state identifier of 40, and the state identifier 0 is the ²Π vibrational ground state without hyperfine splitting.

state identifier (v)	state	HFS
0	² Π v = 0	no
40	² Σ v = 0	no
80	² Π v = 0	yes
81	² Π v = 1	yes